

Enhanced Electroweak Radiative Corrections in SUSY: Gluon-free Observables

I.V.Gaidaenko, A.V.Novikov, V.A.Novikov

ITEP, Moscow, Russia

A.N.Rozanov

ITEP and CPPM, IN2P3, CNRS, Marseille, France

M.I.Vysotsky

INFN, Sezione di Ferrara, Italy and ITEP

Abstract

Large top quark mass is responsible for the enhancement of the oblique radiative corrections in SUSY models. We present the analytical formulas for these corrections to the W -boson mass m_W and to $Z l^+ l^-$ coupling constants. The comparison with the result of the Standard Model fit is made.

The precision measurements of the Z -boson parameters, W -boson mass and top-quark mass have demonstrated that a Standard Model perfectly describes this part of Physics [1]. One has no hopes to improve a Standard Model fit by introducing New Physics since it is already perfect. For the data set announced at HEP97 Conference we obtained $\chi^2/d.o.f. = 18/14$ [2].

Nevertheless, the people still believe that there should be physics beyond the Standard Model and the most popular candidate is supersymmetry. In SUSY models there is a natural reason for the success of the Standard Model in describing the data at the energy scale of the order of the intermediate boson masses. This reason is decoupling: the contribution of SUSY particles to the "low energy" observables is suppressed like $(m_{W,Z}/m_{SUSY})^2$. That is why, when comparing the results of the SUSY model calculations with the experimental data, one gets the lower bounds on superpartner masses. To perform such a program one-loop electroweak radiative corrections in SUSY models should be calculated and this was done, see e.g. [3] - [5]. However, since there are a lot of one-loop diagrams and a large number of parameters even in the simplest MSSM the qualitative picture of electroweak radiative corrections did not emerge. The aim of our investigation is two-fold. Firstly, we will present simple analytical formulas which describe the main part of the radiative corrections in a wide class of SUSY models; secondly, we will include SUSY particles in the description of the electroweak radiative corrections developed and reviewed in [6].

In the present paper we will deal with the observables which are less sensitive to the strong interactions, i.e. with the mass of the W -boson and with $Z l^+ l^-$ -coupling constants g_A and g_V . In future we plan to incorporate Z -boson decays to hadrons which will enable us to perform the general fit of the data and to get lower bounds on the masses of Superpartners.

In order to calculate the radiative corrections to m_W , g_A and g_V one starts with the corrections to G_μ , m_Z and $\bar{\alpha} \equiv \alpha(m_Z)$. The corrections to G_μ are described in SUSY theories by box and vertex diagrams and by the correction to W -boson propagator. The corrections to m_Z and $\bar{\alpha}$ are described by self-energy insertions. Having calculated all the terms one comes to the corrections to m_W , g_A and g_V . The corrections to m_W contain self energy insertions and those to the coupling constants contain the vertices as well. In general all the above corrections are of the order of $(m_{W,Z}/m_{SUSY})^2$ and there are a lot of contributions from a large number of diagrams. However, the violation of $SU(2)_V$ symmetry by large top quark mass penetrates into the SUSY sector of the theory and leads to huge en-

hancement of the corresponding oblique corrections for any value of m_{SUSY} . To begin with, let us neglect $\tilde{t}_L \tilde{t}_R$ mixing. In this case we have with good accuracy: $m_{\tilde{t}_L}^2 - m_{\tilde{b}_L}^2 = m_t^2$ (in what follows we will often designate $\tilde{b}_L$ as $\tilde{b}$). For $(\tilde{t}_L, \tilde{b}_L)$ loop insertion into W -boson propagator we get the result, proportional to $(m_{\tilde{t}_L}^2 - m_{\tilde{b}_L}^2)^2 / m_W^2 m_{SUSY}^2 \approx 16(m_W/m_{SUSY})^2$ since $m_t \approx 2m_W$. That is why the $(\tilde{t}, \tilde{b})$ oblique corrections are responsible for the main part of SUSY corrections to gluon-free observables. (Let us note that the existence of the terms $\sim (m_t)^4$ in the SUSY radiative corrections was observed long ago [7]. Now when we know that top quark is very heavy and from the direct searches it follows that most of SUSY partners should be much heavier than $M_Z/2$ we can claim that the stop-sbottom contribution to the oblique corrections dominates over the other SUSY contributions to gluon-free observables.) The term $\sim m_t^4 / m_W^2 m_{SUSY}^2$ comes from $\Sigma_W(0)$, while the terms proportional to $\Sigma'(0)$ produce the corrections of the order of m_t^2 / m_{SUSY}^2 . Higher order derivatives of self energies are suppressed as $(m_{W,Z}/m_{SUSY})^2$ and should be omitted. Calculating the vector boson self energies and keeping the first two terms in the k^2 expansion, we get the following expressions for the SUSY contributions to the quantities V_i , which are directly related to the physical observables (V_m – to m_W , V_A – to g_A and V_R – to the ratio g_V/g_A , see [6]):

$$\delta_{SUSY} V_A = \frac{1}{m_Z^2} g(m_{\tilde{t}_L}, m_{\tilde{b}}) , \quad (1)$$

$$\delta_{SUSY} V_R = \delta_{SUSY} V_A + \frac{1}{3} Y_L \ln\left(\frac{m_{\tilde{t}_L}^2}{m_{\tilde{b}}^2}\right) , \quad (2)$$

$$\delta_{SUSY} V_m = \delta_{SUSY} V_A + \frac{2}{3} Y_L s^2 \ln\left(\frac{m_{\tilde{t}_L}^2}{m_{\tilde{b}}^2}\right) + \left(\frac{c^2 - s^2}{3}\right) h(m_{\tilde{t}_L}, m_{\tilde{b}}) , \quad (3)$$

where Y_L is the left doublet hypercharge, $Y_L = Q_t + Q_b$, $s \equiv \sin \theta$, $c \equiv \cos \theta$, $\sin^2 \theta \cos^2 \theta = \pi \bar{\alpha}(m_Z) / (\sqrt{2} G_\mu m_Z^2)$ (θ is the electroweak mixing angle) and the functions g and h are standard functions which describe the scalar particle contributions to the vector boson self energies at one loop:

$$g(m_1, m_2) = m_1^2 + m_2^2 - 2 \frac{m_1^2 m_2^2}{m_1^2 - m_2^2} \ln\left(\frac{m_1^2}{m_2^2}\right) , \quad (4)$$

$$h(m_1, m_2) = -\frac{5}{3} + \frac{4m_1^2 m_2^2}{(m_1^2 - m_2^2)^2} + \frac{(m_1^2 + m_2^2)(m_1^4 - 4m_1^2 m_2^2 + m_2^4)}{(m_1^2 - m_2^2)^3} \ln\left(\frac{m_1^2}{m_2^2}\right) . \quad (5)$$

For SUSY partners with masses larger than m_t , $m_{\tilde{b}} \gg m_t$, the leading term in (1) – (3) is universal and originates from $\Sigma_W(0)$ and is given by the function g :

$$\delta_{SUSY} V_i = \frac{1}{3} \frac{(m_{\tilde{t}_L}^2 - m_{\tilde{b}}^2)^2}{m_Z^2 m_{\tilde{b}}^2} . \quad (6)$$

For light SUSY partners $m_{\tilde{b}} \sim m_t$ we should not expand $\delta_{SUSY} V_i$ over $(m_{\tilde{t}_L}^2 - m_{\tilde{b}}^2)/m_{\tilde{b}}^2$ but use expressions (1) – (3) to describe the electroweak corrections. In Figures 1 - 3 the dependences of the functions $\delta_{SUSY} V_i$ on $m_{\tilde{b}}$ are shown. For top quark mass we use $m_t = 175 \text{ GeV}$. The horizontal lines show the difference of the experimental values of V_i and their Standard Model fit. The solid lines correspond to the central values while the dotted ones correspond to the one standard deviation corridor. From the Figures we see that for $m_{\tilde{b}} \geq 200 \text{ GeV}$ SUSY corrections are not essential and the inclusion of SUSY partners will not spoil the Standard Model fit of the data.




Figure 1: The dependence of the function $\delta_{SUSY} V_A$ on $m_{\tilde{b}}$. The horizontal lines show the difference between experimental value of V_A and its fit in the Standard Model, $V_A^{exp} - V_A^{theor} = -0.38(46)$. The solid line corresponds to

the central value, while dotted – to one standard deviation corridor.




Figure 2: The same as Figure 1 for V_R . $V_R^{exp} - V_R^{theor} = -0.0(4)$ was used.




Figure 3: The same as Figure 1 for V_M . $V_m^{exp} - V_m^{theor} = 0.45(65)$ was used.

Now let us take into account $\tilde{t}_L\tilde{t}_R$ mixing which in general is not small being of the order of t -quark mass times the superpartner mass scale, $m_{LR}^2 \sim m_t m_{SUSY}$. As a result of mixing two states with masses m_1 and m_2 are formed, $\tilde{t}_1 = c_u \tilde{t}_L + s_u \tilde{t}_R$ and $\tilde{t}_2 = -s_u \tilde{t}_L + c_u \tilde{t}_R$, where $c_u \equiv \cos \theta_{LR}$, $s_u \equiv \sin \theta_{LR}$ and our functions $\delta_{SUSY}^{LR} V_i$ depend on 4 parameters: m_1 , m_2 , θ_{LR} and $m_{\tilde{b}}$, three of which are independent (the difference of $m_{\tilde{t}_L\tilde{t}_L}^2$ and $m_{\tilde{b}_L\tilde{b}_L}^2$ is determined by the top quark mass). The numerical values of these parameters depend on the SUSY model. Let us present the formulas for the electroweak radiative corrections which take $\tilde{t}_L\tilde{t}_R$ mixing into account:

$$\delta_{SUSY}^{LR} V_A = \frac{1}{m_Z^2} [c_u^2 g(m_1, m_{\tilde{b}}) + s_u^2 g(m_2, m_{\tilde{b}}) - c_u^2 s_u^2 g(m_1, m_2)] \quad , \quad (7)$$

$$\delta_{SUSY}^{LR} V_R = \delta_{SUSY}^{LR} V_A + \frac{1}{3} Y_L [c_u^2 \ln(\frac{m_1^2}{m_{\tilde{b}}^2}) + s_u^2 \ln(\frac{m_2^2}{m_{\tilde{b}}^2})] - \frac{1}{3} c_u^2 s_u^2 h(m_1, m_2) \quad , \quad (8)$$

$$\begin{aligned} \delta_{SUSY}^{LR} V_m &= \delta_{SUSY}^{LR} V_A + \frac{2}{3} Y_L s^2 [c_u^2 \ln(\frac{m_1^2}{m_{\tilde{b}}^2}) + s_u^2 \ln(\frac{m_2^2}{m_{\tilde{b}}^2})] + \\ &+ \frac{c^2 - s^2}{3} [c_u^2 h(m_1, m_{\tilde{b}}) + s_u^2 h(m_2, m_{\tilde{b}})] - \frac{c_u^2 s_u^2}{3} h(m_1, m_2) \quad . \quad (9) \end{aligned}$$

Our approximation should be good for the case when all superpartners have more or less equal masses. When some spartners are considerably lighter than sbottom their contribution can dominate and our approximation can fails.

It is desirable to compare numerically the formulas for the enhanced radiative corrections with the results of full one loop calculations in SUSY models in order to check how good our approximation is.

We are grateful to Z.Berezhiani for discussions on SUSY models. Investigations of I.G., A.N., V.N. and M.V. were supported by RFBR grants 96-02-18010 and 96-15-96578; those of A.N., V.N. and M.V. by INTAS grants 93-3316 and 94-2352 and INTAS-RFBR grant 95-05678 as well. V.A.Novikov is grateful to G.Fiorentini and L.Piemontese for hospitality in Ferrara, where this work was finished.

References

- [1] J.Timmermans, Proceedings of LP 97, Hamburg, 1997;
D.Ward, Proceedings of HEP 97, Jerusalem, 1997;
G.Quast, ibidem.
- [2] M.I.Vysotsky, V.A.Novikov, L.B.Okun and A.N.Rozanov, Standartnaya Model i itogi LEP I, submitted to Yad. Fiz.
- [3] P.Chankowski, A.Dabelstein, W.Hollik, W.Mösle, S.Pokorski and J.Rosiek, Nucl. Phys. **B417** (1994) 101.
- [4] D.Garcia, R.J.Jimenez and J.Sola, Phys. Lett. **B347** (1995) 309, 321.
- [5] W. de Boer, A.Dabelstein, W.Hollik, W.Mösle and U.Schwickerath, Z.Phys. **C75** (1997) 627.
- [6] V.A.Novikov, L.B.Okun, M.I.Vysotsky, Nucl. Phys. **B397** (1993) 35;
V.Novikov, L.Okun, A.Rozanov, M.Vysotsky, LEPTOP, preprint ITEP 19-95/CPPM-95-1; UFN **166** (1996) 539.
- [7] L.Alvarez-Gaume, J.Polchinski and M.Wise, Nucl. Phys. **B211** (1983) 495;
R.Barbieri and L.Maiani, Nucl. Phys. **B224** (1983) 32.